maxill

Instructions For Use

CHLORHEXIDINE GLUCONATE

Purpose: Chlorhexidine gluconate is a broad-spectrum antimicrobial agent used in healthcare and dental for various purposes.

FOR PROFESSIONAL AND INSTITUTIONAL USE ONLY

⚠WARNING!⚠

This product is ***NOT*** to be administered at **FULL** strength. See dilution instructions below.

DANGER
Causes serious eye damage
Provoque des lésions oculaires graves

Wear eye protection/face protection.

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Immediately call a POISON CENTER or doctor/physician. Refer to SDS.

STORAGE: Store at room temperature in a dark place. Do not freeze. Keep out of direct sunlight.

DILUTION INSTRUCTIONS:

Chlorhexidine Gluconate Solution 20% w/v - 473 mL bottle (Item# 40409)

To prepare a 0.12% CHG Rinse, use a suitable Base Rinse such as distilled water, nonionic or cationic mouthwash. Cationic mouthwashes contain cetylpyridinium chloride (E.g. PerioX)

Final Volume of 0.12% CHG Rinse (mL)	=	Volume of Base Rinse (mL)	+	Volume of CHG Concentrate (mL)
1000 mL		994 mL		6 mL
500 mL		497 mL		3 mL
250 mL		248.5 mL		1.5 mL
125 mL		124.25 mL		0.75 mL

Chlorhexidine Gluconate Solution 8.12% w/v - 30 mL bottle (Item# 40416)

To prepare a 0.12% CHG Rinse, add the entire contents of the 30 mL bottle of 8.12% Chlorhexidine Gluconate solution into 2 litres of a Chlorhexidine Gluconate compatible base liquid (e.g. PerioX). Gently agitate the mixture to yield a uniform solution of 0.12% w/v of Chlorhexidine Gluconate.

Spaulding's Classifications

maxill inc.

Canadian Head Office
80 Elm Street
St. Thomas, ON N5R 6C8
Toll Free: 1-800-268-8633
Phone: 1-519-631-7370
Fax: 519-631-3388

USA Head Office
500 West Main Street
Cortland, OH 44410
Toll Free: 1-855-462-9455
Phone: 1-330-637-1176
Fax: 1-330-637-1154

Rev.0: 2025-01-22
MP-REF-40409D